



Medicated Products

YOUR PARTNER
IN VETERINARY
THERAPY

SOLUBLE MEDICINES

MEDICINAL SPECIALTIES

LIQUID MEDICINES

MEDICATED PREMIXES

Amoxifarma

Water soluble powder
for veterinary use

COMPOSITION: 1 kg contains:
Amoxicillin trihydrate 800 g

DIRECTIONS FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Broilers, turkeys, swine

USE AND DOSES:
BROILERS, TURKEYS and SWINE: 0,1-
0,2 g/10 kg b.w. per day (8-16 mg of
amoxicillin trihydrate/kg b.w. per day)

TREATMENT PERIOD:
BROILERS: 3-5 days
TURKEYS: 3-5 days
SWINE: 3-5 days

WITHDRAWAL PERIOD:
BROILERS: 1 days
TURKEYS: 2 days
SWINE: 1 days

THERAPEUTIC INDICATIONS:
Infections sustained by Gram-positive
and Gram-negative bacteria sensi-
tive to amoxicillin: colibacillosis,
Salmonellosis, Staphylococcosis and
Streptococcosis, Red disease and MAM
syndrome of swine, infections of the
Respiratory tract, gastroenteric tract,
urinary tract, genital tract, skin, and
joints; clostridiosis by *Clostridium*
perfringens.

Packing:
143 g plastic bucket
500 g sachet
1,430 g plastic bucket

Clortetraciclina 20%

Water soluble powder
for veterinary use

COMPOSITION: 1 kg contains:
Chlortetracycline 200 g

DIRECTION FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Broilers, layers, swine, sucking calves

USE AND DOSES:
BROILERS: 12,5-25 g/100 kg of body weight (equal to 25-50 mg of chlortetracycline/kg b.w.)

LAYERS: 12,5-25 g/100 kg of body weight (equal to 25-50 mg of chlortetracycline/kg b.w.)

SWINE: 10-25 g/100 kg of body weight (equal to 20-50 mg of chlortetracycline/kg b.w.)

SUCKING CALVES: 12,5-25 g/100 kg of body weight (equal to 25-50 mg of chlortetracycline/kg b.w.)

TREATMENT PERIOD:
BROILERS: 3-5 days
LAYERS: 3-5 days
SWINE: 3-5 days
SUCKING CALVES: 3-5 days

WITHDRAWAL PERIOD:
BROILERS: 6 days
LAYERS: 5 days (eggs)
SWINE: 4 days
SUCKING CALVES: 24 days

THERAPEUTIC INDICATIONS:
BROILERS, LAYERS: Bacterial Diseases of the Respiratory, gastro-intestinal, genital and locomotory tracts, bacterial septicaemia

SWINE: Respiratory and intestinal bacterial Diseases

SUCKING CALVES: Pasteurellosis, infections complicating the viroses of digestive tract, Enzootic Pneumonia

Packing:
1 kg plastic bucket (6x1 carton box)
5 kg bag

Colistina solfato 12%

Water soluble powder
for veterinary use

COMPOSITION: 1 kg contains:
Colistin sulphate 120 g

DIRECTIONS FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Broilers, turkeys, layers, piglets (up to 35 kg b.w.), SUCKING calves

USE AND DOSES:
BROILERS, TURKEYS: 4,15-5 g/100 kg of body weight (equal to 5-6 mg of colistin sulphate/kg b.w.)

LAYERS: 4,15-5 g/100 kg of body weight (equal to 5-6 mg of colistin sulphate/kg b.w.)

PIGLETS: 4,15-5 g/100 kg of body weight (equal to 5-6 mg of colistin sulphate/kg b.w.)

SUCKING CALVES: 4,15-5 g/100 kg of body weight (equal to 5-6 mg of colistin sulphate/kg b.w.)

TREATMENT PERIOD:
BROILERS TURKEYS: 3-15 days
LAYERS: 3-15 days
PIGLETS: 3-15 days
SUCKING CALVES: 3-15 days

WITHDRAWAL PERIOD:
BROILERS, TURKEYS: 2 days
LAYERS: 5 days (eggs)
PIGLETS: 7 days
SUCKING CALVES: 7 days

THERAPEUTIC INDICATIONS:
Bacterial Enteritis due to gram-negative germs (*Salmonella spp*, *Escherichia coli*, *Haemophilus*, *Pasteurella spp*, *Klebsiella spp*, *Shigella spp*, *Pseudomonas*, *Enterobacter spp* e *Proteus spp*)

Packing:
1 kg plastic bucket (6x1 carton box)
5 kg bag

Doxiciclina 50%

Water soluble powder
for veterinary use

COMPOSITION: 1 kg contains:
Doxycycline 500 g

DIRECTION FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Broilers, turkeys, swine

USE AND DOSES:
BROILERS: 4,0 g of product/100 kg b.w.
(20 mg of doxycycline/kg b.w.), cor-
responding to about 15-30 g of prod-
uct/100 litres of water

TURKEYS: 4,0 g of product/100 kg b.w.
(20 mg of doxycycline/kg b.w.), cor-
responding to about 20-40 g of prod-
uct/100 litres of water

SWINE: 2,0 g of product/100 kg b.w.
(equal to 10 mg of doxycycline/kg
b.w.)

TREATMENT PERIOD:
BROILERS: 3-5 days
TURKEYS: 3-5 days
SWINE: 3-5 days

WITHDRAWAL PERIOD:
BROILERS: 6 days
TURKEYS: 8 days
SWINE: 5 days

THERAPEUTIC INDICATIONS:
BROILERS, TURKEYS: Chronic Respira-
tory disease and other Respiratory
illnesses, specially produced by *Pas-
teurella spp*, *Mycoplasma spp*, *Haem-
philus gallinarum*, *Bordetella avium*
and *Chlamydia psittaci*

SWINE: Atrophic rhinitis and Respirato-
ry illnesses specially caused by *Actino-
bacillus spp*, *Bordetella bronchisep-
tica*, *Mycoplasma spp*, *Pasteurella
multocida*, *Streptococcus suis*

Packing:
50 g sachet (50x1 carton box)
1 kg plastic bucket (6x1 carton box)

Eritromicina 15%

Water soluble powder
for veterinary use

COMPOSITION: 1 kg contains:
Erythromycin 150 g

DIRECTION FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Broilers, turkeys, layers

USE AND DOSES:

BROILERS: 13,5-27 g/100 kg of body weight (equal to 20-40 mg of erythromycin/kg b.w.)

TURKEYS: 13,5-27 g/100 kg of body weight (equal to 20-40 mg of erythromycin/kg b.w.)

LAYERS: 6,7-13,5 g/100 kg of body weight (equal to 10-20 mg of erythromycin/kg b.w.)

TREATMENT PERIOD:
BROILERS: 3-5 days
TURKEYS: 3-5 days
LAYERS: 3-5 days

WITHDRAWAL PERIOD:
BROILERS: 3 days
TURKEYS: 6 days
LAYERS: 5 days (eggs)

THERAPEUTIC INDICATIONS:
Mycoplasmosis

Packing:
1 kg plastic bucket (6x1 carton box)
5 kg bag

Eritromicina 20%

Water soluble powder
for veterinary use

COMPOSITION: 1 kg contains:
Erythromycin 200 g

DIRECTION FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Broilers, turkeys, layers

USE AND DOSES:
BROILERS: 10-20 g/100 kg of body weight (equal to 20-40 mg of erythromycin/kg b.w.)

TURKEYS: 10-20 g/100 kg of body weight (equal to 20-40 mg of erythromycin/kg b.w.)

LAYERS: 5-10 g/100 kg of body weight (equal to 10-20 mg of erythromycin/kg b.w.)

TREATMENT PERIOD:
BROILERS: 3-5 days
TURKEYS: 3-5 days
LAYERS: 3-5 days

WITHDRAWAL PERIOD:
BROILERS: 3 days
TURKEYS: 6 days
LAYERS: 5 days (eggs)

THERAPEUTIC INDICATIONS:
Mycoplasmosis

Packing:
1 kg plastic bucket (6x1 carton box)
5 kg bag

Flumechina 20%

Water soluble powder
for veterinary use

COMPOSITION: 1 kg contains:
Flumequine 200 g

DIRECTIONS FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Broilers, swine

USE AND DOSES:
BROILERS: 6,0 g/100 kg of body weight
(equal to 12 mg of flumequine/kg b.w.)

SWINE: 6,0 g/100 kg of body weight
(equal to 12 mg of flumequine/kg b.w.)

TREATMENT PERIOD:
BROILERS: 3-5 days
SWINE: 3-5 days

WITHDRAWAL PERIOD:
BROILERS: 5 days
SWINE: 2 days

THERAPEUTIC INDICATIONS:
BROILERS: Colibacillosis, Salmonellosis, Pasteurellosis

SWINE: Colibacillosis, Salmonellosis, infections sustained by microorganisms sensitive to flumequine

Packing:
1 kg plastic bucket (6x1 carton box)
5 kg bag

Levamisole 30%

Water soluble powder
for veterinary use

COMPOSITION: 1 kg contains:
Levamisole 300 g

DIRECTION FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Swine

USE AND DOSES:
SWINE under 20 kg of b.w.: 2,5-2,6 g of Levamisole 30%/100 kg b.w. (equal to 7,5-8 mg of active principle/kg b.w.)

SWINE from 20 to 40 kg of b.w.: 1,67 g of Levamisole 30%/100 kg b.w. (equal to 5 mg of active principle/kg b.w.)

SWINE over 40 kg of b.w.: 1,35 g of Levamisole 30%/100 kg b.w. (equal to 4 mg of active principle/kg b.w.)

REMARK: in subjects over 190 kg do not exceed 2,5 g of Levamisole 30%/head (equal to 750 mg of levamisole/head)

TREATMENT PERIOD:
SWINE: 1 day. It is advisable to repeat the treatment 14-21 days after the first one

WITHDRAWAL PERIOD:
SWINE: 7 days

THERAPEUTIC INDICATIONS:
Swine infestations caused by adult and larval forms of *Ascarius suum*, *Hyostrogylus spp*, *Metastrongylus spp*, and *Oesophagostomum spp*

Packing:
50 g sachet
1 kg plastic bucket (6x1 carton box)
5 kg bag

Lincomicina 11%

Water soluble powder
for veterinary use

COMPOSITION: 1 kg contains:
Lincomycin..... 110 g

DIRECTIONS FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Swine

USE AND DOSES:
Enteritis: 4,5-9,0 g/100 kg of body weight (equal to 5-10 mg of lincomycin/kg b.w.)

Enzootic Pneumonia: 9,0-18 g/100 kg of body weight (equal to 10-20 mg of lincomycin/kg b.w.)

TREATMENT PERIOD:
SWINE: 7 days

WITHDRAWAL PERIOD:
SWINE: 5 days

THERAPEUTIC INDICATIONS:
Enzootic Pneumonia, Superficial Necrotic Enteritis, Proliferative Enteritis (Ileitis)

Packing:
1 kg plastic bucket (6x1 carton box)
5 kg bag

Ossitetraciclina 20%

Water soluble powder
for veterinary use

COMPOSITION: 1 kg contains:
Oxytetracycline 200 g

DIRECTION FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Broilers, turkeys, layers, swine, SUCK-
ING calves, rabbits

USE AND DOSES:
BROILERS: 20-35 g/100 kg of body
weight (equal to 40-70 mg of oxytetra-
cycline/kg b.w.)

TURKEYS: 20-35 g/100 kg of body
weight (equal to 40-70 mg of oxytetra-
cycline/kg b.w.)

LAYERS: 12,5-25 g/100 kg of body
weight (equal to 25-50 mg of oxytetra-
cycline/kg b.w.)

SWINE: 10-25 g/100 kg of body weight
(equal to 20-50 mg of oxytetracycline/
kg b.w.)

SUCKING CALVES: 12,5-20 g/100 kg of
body weight (equal to 25-40 mg of oxy-
tetracycline/kg b.w.)

RABBITS: 20- 40 g/100 kg of body
weight (equal to 40- 80 mg of oxytet-
racycline/kg b.w.)

TREATMENT PERIOD:
BROILERS: 3-5 days
TURKEYS: 3-5 days
LAYERS: 3-5 days
SWINE: 3-5 days
SUCKING CALVES: 3-5 days
RABBITS: 3-5 days

WITHDRAWAL PERIOD:
BROILERS: 5 days
TURKEYS: 12 days
LAYERS: 5 days (eggs)
SWINE: 9 days
SUCKING CALVES: 10 days
RABBITS: 12 days

THERAPEUTIC INDICATIONS:
BROILERS, TURKEYS, LAYERS: Respira-
tory, gastro-intestinal, genital bacte-
rial diseases, bacterial septicaemia

SWINE: Respiratory and intestinal bac-
terial diseases

SUCKING CALVES: Pasteurellosis, infec-
tions complicating the viroses of diges-
tive tract, Enzootic Pneumonia

RABBITS: Colibacillosis, Salmonellosis,
Pasteurellosis

Packing:
1 kg plastic bucket (6x1 carton box)
5 kg bag

Spectinomycina 50%

Water soluble powder
for veterinary use

COMPOSITION: 1 kg contains:
Spectinomycine 500 g

DIRECTIONS FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Broiler, swine

USE AND DOSES:
BROILER: 50 g/100 litres of drinking
water (equal to 25-50 mg of spectino-
mycine/kg b.w.)

SWINE: 10-16 mg/kg of body weight
(equal to 5-8 mg of spectinomycine/
kg b.w.)

TREATMENT PERIOD:
BROILER: 7 days
SWINE: 3-5 days

WITHDRAWAL PERIOD:
BROILER: 11 days
SWINE: 20 days

THERAPEUTIC INDICATIONS:
BROILER: Respiratory Chronic Diseases

SWINE: Bacterial Enteritis

Packing:
1 kg plastic bucket (6x1 carton box)
5 kg bag

Spiramicina 20%

Water soluble powder
for veterinary use

COMPOSITION: 1 kg contains:
Spiramycine..... 820.000.000 I.U.
(equal to about 200 g)

DIRECTIONS FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Broilers, swine, SUCKING calves

USE AND DOSES:
BROILERS: 250-500 g/100 litres of
drinking water (equal to 0,5-1,0 g of
spiramicine/litre)

SWINE: 12,5 g/100 kg of body weight
(equal to 25 mg of spiramicine/kg
b.w.)

SUCKING CALVES: 12,5 g/100 kg
of body weight (equal to 25 mg of
spiramicine/kg b.w.)

TREATMENT PERIOD:
BROILERS: 3-5 days
SWINE: 3-5 days
SUCKING CALVES: 3-5 days

WITHDRAWAL PERIOD:
BROILERS: 15 days
SWINE: 10 days
SUCKING CALVES: 24 days

THERAPEUTIC INDICATIONS:
BROILERS: Bacterial Diseases of the
Respiratory, gastro intestinal, genital
and locomotory tracts, septicaemia
caused by gram-positive germs

SWINE: Bacterial Enteritis caused by
gram-positive germs, Enzootic Pneu-
monia

SUCKING CALVES: Respiratory Diseases
and neonatal Diseases caused by gram
positive germs

Packing:
1 kg plastic bucket (6x1 carton box)
5 kg bag

Sulfaclopirazina 12,5%

Water soluble powder
for veterinary use

COMPOSITION: 1 kg contains:
Sulphachloropyrazine 125 g

DIRECTION FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Poultry

USE AND DOSES:
POULTRY: 40-60 g/100 kg of body weight (equal to 50-75 mg of sulphachloropyrazine/kg b.w.)

TREATMENT PERIOD:
POULTRY: 5 days

WITHDRAWAL PERIOD:
POULTRY: 10 days

THERAPEUTIC INDICATIONS:
Coccidiosis

Packing:
1 kg plastic bucket (6x1 carton box)
5 kg bag

Sulfadimetossina 20%

Water soluble powder
for veterinary use

COMPOSITION: 1 kg contains:
Sulfadimethoxin..... 200 g

DIRECTIONS FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Broiler, swine (up to 6 months), SUCK-
ING calves, rabbits

USE AND DOSES:
BROILER: 30 g/100 kg of body weight
(equal to 60 mg of sulphadimethoxin/
kg b.w.)

SWINE: 30 g/100 kg of body weight
(equal to 60 mg of sulphadimethoxin/
kg b.w.)

SUCKING CALVES: 12,5-25 g/100 kg of
body weight (equal to 25-50 mg of sul-
phadimethoxin/kg b.w.)

RABBITS: 30 g/100 kg of body weight
(equal to 60 mg of sulphadimethoxin/
kg b.w.)

TREATMENT PERIOD:
BROILER: 3-5 days
SWINE: 3-5 days
SUCKING CALVES: 3-5 days
RABBITS: 3-5 days

WITHDRAWAL PERIOD:
BROILER: 15 days
SWINE: 28 days
SUCKING CALVES: 28 days
RABBITS: 21 days

THERAPEUTIC INDICATIONS:
BROILER, SUCKING CALVES: Bacterial
diseases

SWINE: Bacterial Enteritis

RABBITS: Coccidiosis, Salmonellosis

Packing:
1 kg plastic bucket (6x1 carton box)
5 kg bag

Sulfametazina 20%

Water soluble powder
for veterinary use

COMPOSITION: 1 kg contains:
Sulphamethazine..... 200 g

DIRECTIONS FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Swine (up to 6 months of age)

USE AND DOSES:
SWINE: 75 g/100 kg of body weight
(equal to 150 mg of sulphamethazine/
kg b.w.)

TREATMENT PERIOD:
SWINE: 3-5 days

WITHDRAWAL PERIOD:
SWINE: 28 days

THERAPEUTIC INDICATIONS:
Bacterial Diseases

Packing:
1 kg plastic bucket (6x1 carton box)
5 kg bag

Tiamulina 8%

Water soluble powder
for veterinary use

COMPOSITION: 1 kg contains:
Tiamulin 80 g
(equal to Tiamulin hydrogen fumarate
100 g)

DIRECTION FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Swine

USE AND DOSES:
SWINE: 7,5-11,25 g/100 kg b.w. equal
to about 100-200 g/100 litres of water

TREATMENT PERIOD:
SWINE: 3-5 days

WITHDRAWAL PERIOD:
SWINE: 10 days

THERAPEUTIC INDICATIONS:
Mycoplasmosis, Enzootic Pneumonia,
Superficial Necrotic Enteritis

Packing:
1 kg plastic bucket (6x1 carton box)
5 kg bag

Tilosina 20%

Water soluble powder
for veterinary use

COMPOSITION: 1 kg contains:
Tylosin 200 g

DIRECTION FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Broiler, swine, SUCKING calves

USE AND DOSES:
BROILER: 12,5 g/100 kg of body weight
(equal to 25 mg of tylosin/kg b.w.)

SWINE: 5-6 g/100 kg of body weight
(equal to 10-12 mg of tylosin/kg b.w.)

SUCKING CALVES: 20-25 g/100 kg of
body weight (equal to 40-50 mg of ty-
losin/kg b.w.)

TREATMENT PERIOD:
BROILER: 3-5 days
SWINE: 3-5 days
SUCKING CALVES: 3-7 days

WITHDRAWAL PERIOD:
BROILER: 12 days
SWINE: 8 days
SUCKING CALVES: 24 days

THERAPEUTIC INDICATIONS:
BROILER: Chronic Respiratory Diseases

SWINE: Enzootic Pneumonia, Necrotic
Enteritis

SUCKING CALVES: Pneumonia by
Mycoplasma and/or *Pasteurella Mul-
tocida*

Packing:
1 kg plastic bucket (6x1 carton box)
5 kg bag

Ampifarma

Water soluble powder
for veterinary use

COMPOSITION: 1 kg contains:
Ampicillin 750 g

DIRECTION FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Broilers, swine

USE AND DOSES:
BROILERS: infections of the intestinal tract: 3,3 g of "Ampifarma"/100 kg b.w./day (equal to 2,5 g of ampicillin/100 kg b.w./day), equal to 10-20 g of product/100 litres of water.
In all the other infections: 5,3 g of "Ampifarma"/100 kg b.w./day (equal to 4,0 g of ampicillin/100 kg b.w./day), equal to 30-40 g of product/100 litres of water.

SWINE: 1,3-2,6 g of "Ampifarma"/100 kg b.w. every 12 hours (equal to 1,0-2,0 g of ampicillin/100 kg b.w.)

TREATMENT PERIOD:
BROILERS: 3-5 days
SWINE: 3-5 days

WITHDRAWAL PERIOD:
BROILERS: 4 days
SWINE: 9 days

THERAPEUTIC INDICATIONS:
BROILERS: Infections sustained by *Escherichia coli*, *Salmonellosis*, above all Pullurosis and Typhoid fever, Necrotic Enteritis and gangrenous dermatitis caused by *Clostridium spp.*, *Pasteurellosis*, primary and secondary Staphylococcosis, infective Coryza, secondary bacterial infections, viral and parasitic diseases

SWINE: Colibacillosis, Salmonellosis, Staphylococcosis and Streptococcosis, Red disease and MAM syndrome of swine, infections of the Respiratory tract and gastroenteric tract

Packing:
1.200 g plastic bucket

Chemisole 20%

Water soluble powder
for veterinary use

COMPOSITION: 1 liter contains:
Levamisole 200 g

DIRECTIONS FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Broilers, turkeys

USE AND DOSES:
BROILERS, TURKEYS: 1 ml of "Chemisole 20%" / 8-10 kg b.w. (equal to 20-25 mg of levamisole/kg b.w.), equal to about 50-100 ml / 100 litres of water.

TREATMENT PERIOD:
BROILERS, TURKEYS: 1 day (12 hours)
It is advisable to repeat the treatment 14-21 days after the first one

WITHDRAWAL PERIOD:
BROILERS, TURKEYS: 7 days

THERAPEUTIC INDICATIONS:
Poultry infestations caused by adult and larval forms of *Heterakis gallinarum*, *Capillaria spp*, *Syngamus trachea* and *Ascaridia galli*

Packing:
1 liter plastic bottle (12x1 carton box)
5 liters tank

Chemitrim

Water soluble powder
for veterinary use

COMPOSITION: 1 liter contains:

Trimethoprim 40 g
Sulphadimetoxin 200 g

DIRECTION FOR USE:

Dissolve thoroughly in drinking water

ANIMAL SPECIES:

Swine

USE AND DOSES:

SWINE: 4,5-9,0 ml/100 kg b.w. (0,9-1,8 g of sulfadimethoxine + 0,18-0,36 g of trimethoprim/100 kg b.w.) equal to about 18,0- 45,0 ml/100 litres of drinking water

TREATMENT PERIOD:

SWINE: 3-5 days

WITHDRAWAL PERIOD:

SWINE: 7 days

THERAPEUTIC INDICATIONS:

Bacterial infections, especially those caused by Coliforms and Salmonellae, atrophic rhinitis and Respiratory Diseases

Packing:

1 liter plastic bottle (12x1 carton box)
5 liters tank

TAF 25%

Water soluble powder
for veterinary use

COMPOSITION: 1 liter contains:
Thiamphenicol..... 250 g

DIRECTION FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Broilers, turkeys

USE AND DOSES:
BROILERS and TURKEYS: 1,6-2,0 ml
of TAF 25%/10 kg b.w. (equivalent to
40-50 mg of thiamphenicol/kg b.w.),
equal to about 80-125 ml/100 litres of
water

TREATMENT PERIOD:
BROILERS: 3-5 days
TURKEYS: 3-5 days

WITHDRAWAL PERIOD:
BROILERS: 4 days
TURKEYS: 6 days

THERAPEUTIC INDICATIONS:
BROILERS: Bacterial diseases of the
respiratory and intestinal system,
therapy of Chronic Respiratory disease
and in avian Coryza caused especially
by Gram-positive and Gram-negative
micro-organisms, Rickettsiae, Myco-
plasmas and Clamydiae.

TURKEYS: bacterial infections sub-
tained by *Salmonella spp*, *Pasteurella
multocida*, *Ornithobacterium rhi-
notracheale*, *Staphylococcus aureus*,
Clostridium perfringens, *Mycoplasma
gallisepticum*, *Mycoplasma synoviae*,
Mycoplasma meleagridis.

Packing:
1 liter plastic bottle (12x1 carton box)
5 liters tank

Amprosid

Oral solution
for veterinary use

COMPOSITION: 1 liter contains:
Amprolium HCl 250 g

DIRECTIONS FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Broilers, layers

USE AND DOSES:
BROILERS: 200 ml/ 100 Litres of drinking water (equal to 6-35 mg of amprolium Hcl/kg b.w.)

LAYERS: 24-100 ml/ 100 Litres of drinking water (equal to 6-35 mg of amprolium Hcl/kg b.w.)

TREATMENT PERIOD:
BROILERS: 7 days
LAYERS: 7 days

WITHDRAWAL PERIOD:
BROILERS: 0 days
LAYERS: 0 days (eggs)

THERAPEUTIC INDICATIONS:
Coccidiosis

Packing:
1 liter plastic bottle (12x1 carton box)
5 liters tank

Colisid

Oral solution
for veterinary use

COMPOSITION: 1 liter contains:
Colistin sulphate 120 g

DIRECTIONS FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Broilers, turkeys, layers, piglets (up to 35 kg b.w.), calves, rabbits

USE AND DOSES:
BROILERS, TURKEYS, LAYERS: 4,2-5 ml/100 kg of b.w. (equal to 5 -6 mg of colistin sulphate/kg b.w.)

PIGLETS: 4,2-5 ml/100 kg of b.w. (equal to 5-6 mg of colistin sulphate/kg b.w.)

CALVES: 4,2-5 ml/100 kg of b.w. (equal to 5-6 mg of colistin sulphate/kg b.w.)

RABBITS: 4,2-5 ml/100 kg of b.w. (equal to 5-6 mg of colistin sulphate/kg b.w.)

TREATMENT PERIOD:
BROILERS, TURKEYS, LAYERS: 3-15 days
PIGLETS: 3-15 days
CALVES: 3-15 days
RABBITS: 3-15 days

WITHDRAWAL PERIOD:
Broilers: 0 days
Turkeys: 1 day
Layers: 0 days (eggs)
Rabbits: 5 days
Piglets: 7 days
Calves: 7 days

THERAPEUTIC INDICATIONS:
Bacterial Enteritis due to gram-negative germs (*Salmonella spp*, *Escherichia coli*, *Haemophilus*, *Pasteurella spp*, *Klebsiella spp*, *Shigella spp*, *Pseudomonas*, *Enterobacter spp* e *Proteus spp*)

Packing:
1 liter plastic bottle (12x1 carton box)
5 liters tank

Flumechina 50%

Oral solution
for veterinary use

COMPOSITION: 1 liter
Flumequine 500 g

DIRECTIONS FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Broilers

USE AND DOSES:
BROILERS: 2,4 g/100 g of b.w. (equal
to 12 mg of flumequine/kg of b.w.)

TREATMENT PERIOD:
BROILERS: 3-5 days

WITHDRAWAL PERIOD:
BROILERS: 2 days

THERAPEUTIC INDICATIONS:
Colibacillosis, Salmonellosis, Pasteurel-
losis

Packing:
1 liter plastic bottle (12 x 1 carton box)
5 liters tank

Ossitetraciclina 20%

Oral solution
for veterinary use

COMPOSITION: 1 liter contains:
Oxytetracycline 200 g

DIRECTIONS FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Broilers, turkeys, layers, swine,
calves, rabbits

USE AND DOSES:
BROILERS: 20-35 g/100 kg of body
weight (equal to 40-70 mg of oxytetra-
cycline/kg b.w.)

TURKEYS: da 20 a 35 g/100 kg of body
weight (equal to 40-70 mg of oxytetra-
cycline/kg b.w.)

LAYERS: 12,5 a 25 g/100 kg of body
weight (equal to 25-50 mg of oxytetra-
cycline/kg b.w.)

SWINE: 10-25 g/100 kg of body weight
(equal to 20-50 mg of oxytetracycline/
kg b.w.)

CALVES: 12,5-20 g/100 kg of body
weight (equal to 25-40 mg of oxytetra-
cycline/kg b.w.)

RABBITS: 20-40 g/100 kg of body
weight (equal to 40-80 mg of oxytetra-
cycline/kg b.w.)

TREATMENT PERIOD:

BROILERS: 3-5 days
TURKEYS: 3-5 days
LAYERS: 3-5 days
SWINE: 3-5 days
CALVES: 3-5 days
RABBITS: 3-5 days

WITHDRAWAL PERIOD:

BROILERS: 5 days
TURKEYS: 12 days
LAYERS: 5 days (eggs)
SWINE: 18 days
CALVES: 20 days
RABBITS: 12 days

THERAPEUTIC INDICATIONS:

BROILERS, TURKEYS, LAYERS: Bacte-
rial Diseases of the Respiratory gastro-
intestinal and genital tracts, bacterial
septicaemia

SWINE: Respiratory and intestinal bac-
terial Diseases

CALVES: Pasteurellosis, infections
complicating the viroses of digestive
tract, Enzootic Pneumonia

RABBITS: Colibacillosis, Salmonellosis
and Pasteurellosis

Packing:

1 liter plastic bottle (12x1 carton box)
5 liters tank

Spiramicina 20%

Oral solution
for veterinary use

COMPOSITION: 1 ml contains:
Spiramycine.....820.000 I.U.

DIRECTIONS FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Broilers, swine

USE AND DOSES:
BROILERS: 250-500 ml/100 litres of
drinking water (equal to 287.000 I.U.
of spiramycine/kg b.w.)

SWINE: 12,5 ml/100 kg of body
weight equal (equal to 102.500 I.U. of
spiramycine/kg b.w.)

TREATMENT PERIOD:
BROILERS: 3-5 days
SWINE: 3-5 days

WITHDRAWAL PERIOD:
BROILERS: 10 days
SWINE: 20 days

THERAPEUTIC INDICATIONS:
BROILERS: Bacterial Diseases of the
Respiratory, gastro-intestinal, genital
and locomotory tracts, septicaemia
caused by gram-positive germs

SWINE: Bacterial Enteritis by gram-
positive germs, Enzootic Pneumonia

Packing:
1 liter plastic bottle (12x1 carton box)
5 liters tank

Sulfachinossalina 20%

Oral solution
for veterinary use

COMPOSITION: 1 liter contains:
Sulphaquinoxaline 200 g

DIRECTIONS FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Broilers, rabbits

USE AND DOSES:
BROILERS: 30 g/100 kg of body weight
(equal to 60 mg of sulphaquinoxaline/
kg b.w.)

RABBITS: 30 g/100 kg of body weight
(equal to 60 mg of sulphaquinoxaline/
kg b.w.)

TREATMENT PERIOD:
BROILERS: 3-5 days
RABBITS: 3-5 days

WITHDRAWAL PERIOD:
BROILERS: 35 days
RABBITS: 22 days

THERAPEUTIC INDICATIONS:
BROILERS: Bacterial Enteritis, Coccidi-
osis

RABBITS: Coccidiosis, Salmonellosis

Packing:
1 liter plastic bottle (12x1 carton box)
5 liters tank

Tiamulina 10%

Oral solution
for veterinary use

COMPOSITION: 1 liter contains:
Tiamulin 100 g
(equal to Tiamulin hydrogen fumarate
125 g)

DIRECTIONS FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Broilers, swine

USE AND DOSES:
BROILERS: 125-200 g/100 litres of
drinking water (equal to 20-40 mg of
tiamulin/kg b.w.)

SWINE: 5-10 g/100 kg b.w. equal to
about 100-200 g of product/100 litres
of drinking water of liquid feed

TREATMENT PERIOD:
BROILERS: 3-5 days
SWINE: 5 days

WITHDRAWAL PERIOD:
BROILERS: 5 days
SWINE: 4 days

THERAPEUTIC INDICATIONS:
BROILERS: Mycoplasmosis Chronic Res-
piratory Diseases, infectious Arthritis

SWINE: Mycoplasmosis, Enzootical
Pneumonia, necrotical Enteritis (swine
dysentery) sustained by bacteria sensi-
tive to tiamulin

Packing:
1 liter plastic bottle (12x1 carton box)
5 liters tank

Tilosina 20%

Oral solution
for veterinary use

COMPOSITION: 1 liter contains:
Tylosin 200 g

DIRECTIONS FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Broilers, swine, sucking calves

USE AND DOSES:
BROILERS: 12,5 g/100 kg b.w. equal to
25 mg of tylosin/kg of b.w.

SWINE: 5-6 g/100 kg b.w. (equal to 10-
12 mg of tylosin/kg b.w.)

SUCKING CALVES: 20-25 g/100 kg b.w.
(equal to 40-50 mg of tylosin/kg b.w.)

TREATMENT PERIOD:
BROILERS: 3-5 days
SWINE: 3-7 days
SUCKING CALVES: 3-7 days

WITHDRAWAL PERIOD:
BROILERS: 1 day
SWINE: 8 days
SUCKING CALVES: 24 days

THERAPEUTIC INDICATIONS:
BROILERS: Chronic Respiratory Dis-
eases

SWINE: Enzootical Pneumonia and
Necrotic Enteritis

SUCKING CALVES: Pneumonia by *Mycoplasma* and/or *Pasteurella spp*

Packing:
1 liter plastic bottle (12x1 carton box)
5 liters tank

Tilosina 24%

Oral solution
for veterinary use

COMPOSITION: 1 liter contains:
Tylosin 200 g
(equal to Tylosin tartrate 240 g)

DIRECTIONS FOR USE:
Dissolve thoroughly in drinking water

ANIMAL SPECIES:
Broilers, turkeys

USE AND DOSES:
BROILERS: 12,5 ml/100 kg b.w. equal
to 2,5 g of tylosin/100 kg of b.w., cor-
responding to 50-100 ml/100 litres of
drinking water

TURKEYS: 12,5 ml/100 kg b.w. equal
to 2,5 g of tylosin/100 kg of b.w., cor-
responding to 50-100 ml/100 litres of
drinking water

TREATMENT PERIOD:
BROILERS: 3-5 days
TURKEYS: 3-5 days

WITHDRAWAL PERIOD:
BROILERS: 2 day
TURKEYS: 2 day

THERAPEUTIC INDICATIONS:
BROILERS, TURKEYS: Chronic Respira-
tory Diseases

Packing:
1 liter plastic bottle (12x1 carton box)
5 liters tank

Trimetoprim 40 Sulfadimetossina 200

Oral solution
for veterinary use

COMPOSITION: 1 liter contains:

Trimethoprim 40 g
Sulphadimetoxin 200 g

DIRECTIONS FOR USE:

Dissolve thoroughly in drinking water

ANIMAL SPECIES:

Rabbits, broilers (layers excluded),
turkeys

USE AND DOSES:

BROILERS: 10-20 g/100 kg b.w., equal
to 2-4 g of sulphadimetoxin and 0,4-
0,8 g of trimethoprim/100 kg of b.w.

TURKEYS: 10-20 g/100 kg b.w., equal
to 2-4 g of sulphadimetoxin and 0,4-
0,8 g of trimethoprim/100 kg of b.w.

RABBITS: 10-20 g/100 kg b.w., equal to
2-4 g of sulphadimetoxin + 0,4-0,8 g of
trimethoprim/100 kg of b.w.

TREATMENT PERIOD:

RABBITS: 5-6 days
BROILERS: 5-6 days
TURKEYS: 5-6 days

WITHDRAWAL PERIOD:

BROILERS: 8 days
TURKEYS: 7 days
RABBITS: 11 days

THERAPEUTIC INDICATIONS:

Respiratory Diseases and urogenital in-
fections

Packing:

1 liter plastic bottle (12x1 carton box)
5 liters tank

Anprociclina 200

Premix for
veterinary use only

COMPOSITION: 1 kg contains:
Oxytetraciline..... 200 g

DIRECTIONS FOR USE:
Mix carefully in feed

ANIMAL SPECIES:
Swine, rabbits, fish

USE AND DOSES:
SWINE: 200-1250 g/100 Kg of feed
(equal to 20-50 mg of oxytetracycline/
kg b.w.)

RABBITS: 275-770 g/100 Kg of feed
(equal to 40-80 mg of oxytetracycline/
kg b.w.)

FISH: 1750-3750 g/100 Kg of feed
(equal to 75 mg of oxytetracycline/kg
b.w.)

TREATMENT PERIOD:

SWINE: 3-5 days
RABBITS: 3-5 days
FISH: 7-10 days

WITHDRAWAL PERIOD:

SWINE: 14 days
RABBITS: 12 days
FISH: 37 days

THERAPEUTIC INDICATIONS:

SWINE: Respiratory and intestinal bacterial Diseases caused by *Actinobacillus pleuro Pneumoniae*, *Bordetella bronchiseptica*, *Brucella spp*, *Clostridium spp*, *Erysipelothrix Rhusiopatiae*, *Mycobacterium*, *Mycoplasma spp*, *Pasteurella multocida*, *Pseudomonas spp*, *Streptococcus spp*

RABBITS: Colibacillosis, Salmonellosis, Pasteurellosis

FISH: Bacterial Diseases caused by *Aeromonas spp*, *Cytophaga*, *Pseudomonas spp*, *Vibrio Anguillarum*

Packing:
25 kg bag

Anprometazina 200

Premix for
veterinary use only

COMPOSITION: 1 kg contains:
Sulphamethazine..... 200 g

DIRECTIONS FOR USE:
Mix carefully in feed

ANIMAL SPECIES:
Swine (up to 6 months)

USE AND DOSES:
SWINE: 1500-3750 g/100 Kg of feed
(equal to 150 mg of sulphamethazine/
kg b.w.)

TREATMENT PERIOD:
SWINE: 3-5 days

WITHDRAWAL PERIOD:
SWINE: 28 days

THERAPEUTIC INDICATIONS:
Bacterial Diseases

Packing:
25 kg bag

Anprotiazolo 200

Premix for
veterinary use only

COMPOSITION: 1 kg contains:
Formylsulphathiazole 200 g

DIRECTIONS FOR USE:
Mix carefully in feed

ANIMAL SPECIES:
Swine (up to 6 months of age), rabbits

USE AND DOSES:
SWINE: 1800-2500 g/100 Kg of feed
(equal to 150-200 mg of formylsulphathiazole/kg b.w.)

RABBITS: 800-1000 g/100 Kg of feed
(equal to 150-250 mg of formylsulphathiazole/kg b.w.)

TREATMENT PERIOD:
SWINE: 3-5 days
RABBITS: 7 days

WITHDRAWAL PERIOD:
SWINE: 7 days
RABBITS: 5 days

THERAPEUTIC INDICATIONS:
Bacterial Enteritis

Packing:
25 kg bag

Anpro Spiramix 200

Premix for
veterinary use only

COMPOSITION: 1 kg contains:
Spiramycine..... 200 g
(equal to at least 6.400.000.000 I.U.)

DIRECTIONS FOR USE:
Mix carefully in feed

ANIMAL SPECIES:
Broilers, swine

USE AND DOSES:
BROILERS: 250-500 g/100 Kg of feed
(equal to 70 mg of spiramycine/kg
b.w.)

SWINE: 200-500 g/100 Kg of feed
(equal to 25 mg of spiramycine/kg
b.w.)

TREATMENT PERIOD:
BROILERS:3-5 days
SWINE:3-5 days

WITHDRAWAL PERIOD:
BROILERS: 15 days
SWINE: 20 days

THERAPEUTIC INDICATIONS:
BROILERS: Bacterial Diseases of the
Respiratory, gastro intestinal and loco-
motory tracts, Septicaemia caused by
gram-positive germs

SWINE: Bacterial Enteritis caused by
gram-positive germs, Enzootic Pneu-
monia

Packing:
25 kg bag

Premix for
veterinary use only

Chemisole 30% PX

COMPOSITION: 1 kg contains:
Levamisole 300 g

DIRECTIONS FOR USE:
Mix carefully in feed

ANIMAL SPECIES:
Swine

USE AND DOSES:
SWINE under 20 kg b.w.: 70-85 g/100 kg of feed (equal to 7,5-8,0 mg of levamisole/kg b.w.)

SWINE from 20 to 40 kg b.w.: 48-60 g/100 kg of feed (equal to 5,0 mg of levamisole/kg b.w.)

SWINE over 40 kg b.w.: 40-55 g/100 kg of feed (equal to 4,0 mg of levamisole/kg b.w.)

REMARK: in subjects over 190 kg b.w. do not exceed 2,5 g/head, equal to 750 mg of levamisole per head

TREATMENT PERIOD:
SWINE: 1 day
It is advisable to repeat the treatment 14-21 days after the first one

WITHDRAWAL PERIOD:
7 days

THERAPEUTIC INDICATIONS:
Infestations caused by adult and larval forms of *Ascaris suum*, *Hyostrongylus spp*, *Metastrongylus spp*, *Oesophagostomum spp*

Packing:
25 kg bag

Clortetraciclina 200

Premix for
veterinary use only

COMPOSITION: 1 kg contains:
Chlortetracycline HCL..... 200 g

DIRECTIONS FOR USE:
Mix carefully in feed

ANIMAL SPECIES:
Broilers, layers, swine

USE AND DOSES:
BROILERS: 75-400 g/100 Kg of feed
(equal to 25-50 mg of chlortetracycline/kg b.w.)

LAYERS: 200-500 g/100 Kg of feed
(equal to 25-50 mg of chlortetracycline/kg b.w.)

SWINE: 200-1250 g/100 Kg of feed
(equal to 20-50 mg of chlortetracycline/kg b.w.)

TREATMENT PERIOD:
BROILERS: 3-5 days
LAYERS: 3-5 days
SWINE: 3-5 days

WITHDRAWAL PERIOD:
BROILERS: 6 days
LAYERS: 9 days (eggs)
SWINE: 9 days

THERAPEUTIC INDICATIONS:
BROILERS, LAYERS: Bacterial Diseases of the respiratory, gastro-intestinal and genital tracts. Bacterial septicaemia

SWINE: Respiratory and intestinal bacterial diseases

Packing:
25 kg bag

Colifarm

Premix for
veterinary use only

COMPOSITION: 1 kg contains:
Flumequine 200 g

DIRECTIONS FOR USE:
Mix carefully in feed

ANIMAL SPECIES:
Fish, swine

USE AND DOSES:
FISH: 600 g/100 Kg of feed (equal to 12 mg of flumequine/kg b.w.)

SWINE: 150-500 g/100 Kg of feed (equal to 12 mg of flumequine/kg b.w.)

TREATMENT PERIOD:
FISH: 5 days
SWINE: 5 days

WITHDRAWAL PERIOD:
FISH: 8 days (70° C/day)
SWINE: 2 days

THERAPEUTIC INDICATIONS:
FISH: Forunculosis from *Aeromonas*,
Vibriosis

SWINE: Colibacillosis, Salmonellosis,
infections due to microorganisms sensitive to flumequine

Packing:
25 kg bag

Dimetossin 200

Premix for
veterinary use only

COMPOSITION: 1 kg contains:
Sulphadimetoxin 200 g

DIRECTIONS FOR USE:
Mix carefully in feed

ANIMAL SPECIES:
Broiler, swine (up to 6 months), rabbits

USE AND DOSES:
BROILER: 200-475 g/100 Kg of feed
(equal to 60 mg of sulphadimetoxin/
kg b.w.)

SWINE: 600-1550 g/100 Kg of feed
(equal to 60 mg of sulphadimetoxin/
kg b.w.)

RABBITS: 400-500 g/100 Kg of feed
(equal to 60 mg of sulphadimetoxin/
kg b.w.)

TREATMENT PERIOD:
BROILER: 3-5 days
SWINE: 3-5 days
RABBITS: 3-5 days

WITHDRAWAL PERIOD:
BROILER: 15 days
SWINE: 28 days
RABBITS: 21 days

THERAPEUTIC INDICATIONS:
BROILER: Bacterial Diseases
SWINE: Bacterial Enteritis
RABBITS: Coccidiosis, Salmonellosis

Packing:
25 kg bag

Premix for
veterinary use only

Erytromycina 200

COMPOSITION: 1 kg contains:
Erythromycin 200 g

DIRECTIONS FOR USE:
Mix carefully in feed

ANIMAL SPECIES:
Broilers, layers

USE AND DOSES:
BROILERS: 75-300 g/100 Kg of feed
(equal to 20-40 mg of erythromycin/
kg b.w.)

LAYERS: 110-250 g/100 Kg of feed
(equal to 10-20 mg of erythromycin/
kg b.w.)

TREATMENT PERIOD:
BROILERS: 3-5 days
LAYERS: 3-5 days

WITHDRAWAL PERIOD:
BROILERS: 6 days
LAYERS: 5 days (eggs)

THERAPEUTIC INDICATIONS:
Mycoplasmosis

Packing:
25 kg bag

Lincofarm 110

Premix for
veterinary use only

COMPOSITION: 1 kg contains:
Lincomycin..... 110 g

DIRECTIONS FOR USE:
Mix carefully in feed

ANIMAL SPECIES:
Swine

USE AND DOSES:
SWINE: 200 g/100 Kg of feed (equal to
10-20 mg of lincomycin/kg b.w.)

TREATMENT PERIOD:
SWINE: 7 days

WITHDRAWAL PERIOD:
SWINE: 1 day

THERAPEUTIC INDICATIONS:
Enzootic Pneumonia, Superficial
Necrotic Enteritis, Proliferative Enteri-
tis (Ileitis) caused by microorganisms
sensitive to lincomycin

Packing:
25 kg bag

Premix for
veterinary use only

Medistin 120

COMPOSITION: 1 kg contains:
Colistin Sulphate 120 g

DIRECTIONS FOR USE:
Mix carefully in feed

ANIMAL SPECIES:
Piglets (up to 35 kg b.w.), broiler, turkeys, layers

USE AND DOSES:
PIGLETS: 100 g/100 Kg of feed (equal to 5,0-6,0 mg of colistin sulphate/kg b.w.)

BROILER: 83 g/100 Kg of feed (equal to 5,0-6,0 mg of colistin sulphate/kg b.w.)

TURKEYS: 83 g/100 Kg of feed (equal to 5,0-6,0 mg of colistin sulphate/kg b.w.)

LAYERS: 83 g/100 Kg of feed (equal to 5,0-6,0 mg of colistin sulphate/kg b.w.)

TREATMENT PERIOD:
PIGLETS: 3-15 days
BROILER: 3-15 days
TURKEYS: 3-15 days
LAYERS: 3-15 days

WITHDRAWAL PERIOD:
PIGLETS: 2 days
BROILER: 5 days
TURKEYS: 5 days
LAYERS: 5 days (eggs)

THERAPEUTIC INDICATIONS:
Bacterial Enteritis sustained by gram-negative germs (*Salmonella spp*, *Escherichia coli*, *Haemophilus*, *Pasteurella spp*, *Klebsiella spp*, *Shigella spp*, *Pseudomonas*, *Enterobacter spp* e *Proteus spp*)

Packing:
25 kg bag

Spectin 500

Premix for
veterinary use only

COMPOSITION: 1 kg contains:
Spectinomycine 500 g

DIRECTIONS FOR USE:
Mix carefully in feed

ANIMAL SPECIES:
Swine

USE AND DOSES:
SWINE: 25-50 g/100 Kg of feed (equal
to 4-8 mg of spectinomycine/kg b.w.)

TREATMENT PERIOD:
SWINE: 3-5 days

WITHDRAWAL PERIOD:
SWINE: 20 days

THERAPEUTIC INDICATIONS:
Bacterial Enteritis

Packing:
25 kg bag

Tiamix

Premix for
veterinary use only

COMPOSITION: 1 kg contains:
Tiamulin 80 g
(equal to tiamulin hydrogen fumarate
100 g)

DIRECTIONS FOR USE:
Mix carefully in feed

ANIMAL SPECIES:
Swine

USE AND DOSES:
SWINE: 156-250 g/100 Kg of feed
(equal to 3,7-6,0 mg of tiamulin/kg
b.w.)

TREATMENT PERIOD:
SWINE: 3-5 days

WITHDRAWAL PERIOD:
SWINE: 4 days

THERAPEUTIC INDICATIONS:
Mycoplasmosis, Enzootic Pneumonia,
Superficial Necrotic Enteritis

WARNING: Do not administer in com-
bination with monensin, narasin,
salinomycin and lasalocid

Packing:
25 kg bag

Tiam-oxi

Premix for
veterinary use only

COMPOSITION: 1 kg contains:

Tiamulin 40 g
(equal to tiamulin hydrogen fumarate
50 g)
Oxytetracine 300 g

DIRECTIONS FOR USE:

Mix carefully in feed

ANIMAL SPECIES:

Swine

USE AND DOSES:

SWINE: 300-500 g/100 Kg of feed
(equal to 45 mg of oxytetracine + 6
mg tiamulin/kg b.w.)

TREATMENT PERIOD:

SWINE: 5 days

WITHDRAWAL PERIOD:

SWINE: 8 days

THERAPEUTIC INDICATIONS:

Mycoplasmosis, complications post-influenza or during pseudorabies, Respiratory Diseases especially Enzootic Pneumonia, infective pleuropneumonia, atrophic rhinitis, intestinal Diseases especially colibacillosis, Superficial Necrotic Enteritis, Salmonellosis, diseases of the motor system and the serous membranes especially arthritis, polyserositis and foot-paronychia

WARNING: Do not administer in combination with monensin, narasin, salinomycin and lasalocid

Packing:
25 kg bag

Premix for
veterinary use only

Tylosin 200 premix

COMPOSITION: 1 kg contains:
Tylosin tartrate..... 200 g

DIRECTIONS FOR USE:
Mix carefully in feed

ANIMAL SPECIES:
Broiler, swine

USE AND DOSES:
BROILER: 150-400 g/100 Kg of feed
(equal to 25 mg of tylosine/kg b.w.)

SWINE: 100-300 g/100 Kg of feed
(equal to 10-12 mg of tylosine/kg b.w.)

TREATMENT PERIOD:
BROILER: 3-5 days
SWINE: 3-5 days

WITHDRAWAL PERIOD:
BROILER: 12 days
SWINE: 16 days

THERAPEUTIC INDICATIONS:
BROILER: Chronic Respiratory Diseases
SWINE: Enzootic Pneumonia, Necrotic
Enteritis

Packing:
25 kg bag

